vin® Insect Killer Lawn Granules will provide control of the following pests at the application ra		
Pest indis (including Greenbug) myworms (including Lawn Armyworm, Fall Armyworm) lbugs (including Bluegrass Billbug, Hunting Billbug) ick Turfgrass Ataenius (Adult) stletail terpillars (including Lucerne Moth) inch Bugs ckets (including House Cricket, Common Short-tailed Cricket, Indies Short-tailed Cricket, Arizona Cricket, Tropical Cricket, Variable Field Cricket, Fall Field Cricket, Southeastern Field Cricket, Texas Field Cricket, Spring Field Cricket, Northern Wood Cricket, Western Striped Cricket, Eastern Striped Cricket, Sand Field Cricket, Vocal Field Cricket, Southern Wood Cricket) tworms (including Black Cutworm, Bronzed Cutworm, Granulate Cutworm, Variegated Cutworm) rwigs ry Skipper ebbrat t Fly asshoppers dybeetles affloppers balybugs lbugs lales (including Ground Pearls) ed Maggots wbugs ringtails iped Grassworm o-lined Spittle Bugs ebworms (including Sod Webworm, Grass Webworm, Tropical Sod Webworm) reworms reworms	1 to 2 lbs per 1,000 ft ²	
nual Bluegrass Weevil (Hyperodes) rmudagrass Mites iggers voer Mites ane Flies (larvae) ound Beetle ound Pearl ider Mites lie Crickets (including Tawny Mole Cricket, Southern Mole Cricket, Short-Winged Mole Cricket) iders (including Brown Recluse Spider, Marbled Cellar Spider, Longbodied Cellar Spider, Shortbodied Cellar Spider, Thinlegged Wolf Spider, Burrowing Wolf Spider, Pirate Wolf Spider) ts (except Carpenter and Pharaoh Ants)	2 to 4 lbs per 1,000 ft ²	OK LAWNYS * BROOMS
Is (except Carpenter and Pilaradii Ants) ropean Chafer as (larvae & adults) ubs ported Fire Ants (adults) (excluding mound treatments) llipedes arab Beetles (includes Japanese Beetle, Green June Beetle, May Beetle) orpions (including Arizona Stripetail Scorpion, Southern Unstriped Scorpion, Lesser Stripetail Scorpion, Yellow Ground Scorpion, California Common Scorpion, Northern Scorpion, Eastern Sand Scorpion) ks (including Brown Dog Tick and ticks that may transmit Lyme Disease)	4 lbs per 1,000 ft ²	
continum control of particular pests, follow recommended procedures outlined below: s: Mound Treatment, including Fire Ants: Treat ant mounds by applying ½ cup of this product per moicient force to break its apex and allow the insecticide to flow into the ant tunnels, but not sufficient to sage the mound that may lead to relocation. For best results, apply in cool weather (65 to 80°F) or in earing hours. Treat new mounds as they appear. This product will provide control of ant mounds within itsils such as teaspoons and measuring cups should not be used for food purposes after use with inset mum control or for heavily infested areas, combine both mound and broadcast treatments. Insymorms, Cutworms, and Sod Webworms: Irrigate the treated area with up to 0.1 inches of water imm control or activate (release from the granule) the insecticide. ual Bluegrass Weevil (Hyperodes) adults: Time application for when adult weevils leave their overwire into grass areas. bug adults: Apply when adult billbugs are first observed during April and May. ck Turfgrass Ataenius adults: Apply during May and July to control the first and second generation of sinus adults. nch Bugs: Irrigate the treated area with up to 0.25 inches of water immediately after application to activate provide in the summer. In temperate regions, Spring applications will provide control of over-wintered a larvae: Irrigate the treated area with up to 0.5 inches of water immediately after application to activate unle) the insecticide. It crickets: Apply as late in the day as possible to moist soil (irrigate before application in necessary), a larvae: Irrigate the treated area with up to 0.5 inches of water immediately after application to activate pulse the insecticide. It crickets: Apply as late in the day as possible to moist soil (irrigate before application in fecessary), a finess: (including ticks that transmit Lyme Disease and Rocky Mountain Spotted fever): Target applications of control larger, more damaging, nymphs later in the year. prijons: Tr	significantly rity morning or late four hours. Food cticide. For nediately after netering sites and black turfgrass rate (release from both nymphs and chinch bugs. e (release from the nd water in with up e crickets in the her and more reside hidden during prior to treatment. ons to kill juvenile mmer and fall areas and around rty boundaries from the granules ided control, do not tion rates when ling areas on host leaf litter. erican dog ticks vious infestation is III. Inspections can in a square foot of r application to	
lies Short-tailed Cricket, Arizona Cricket, Tropical Cricket, Variable Field Cricket, Fall Field Cricket, Socket, Texas Field Cricket, Arizona Cricket, Northern Wood Cricket, Western Striped Cricket, Easte and Field Cricket, Voal Field Cricket, Southern Wood Cricket), Cutworms (including Black Cutworm, Ind Field Cricket, Outworms (including Black Cutworm), Dichondra Flea Beetle, Earwigs, Essex Skipper, European Chane Fly Larvae, Fiery Skipper, Fire Ants, Firebrat, Fleas (larvae and adult), Frit Fly, Grasshoppers, Gree ette, Ground Pearl, Grubs, Ladybeetles (including Convergent Ladybeetle, Seven-spotted Ladybeetle, dybeetle, Thirteen-spotted Ladybeetle, Twice-stabbed Ladybeetle, Asian Ladybeetle, Mealybug Ladybicerne Moth, Mealybugs, Millipedes, Mole Crickets (including Tawny Mole Cricket, Southern Mole Crict-twinged Mole Cricket), Pillbugs, Scarab beetles (including Japanese Beetle, Green June Beetle, Morpions (including Arizona Stripetail Scorpion, Southern Unstriped Scorpion, Lesser Stripetail Scorpior, California Common Scorpion, Northern Scorpion, Eastern Sand Scorpion), Seed Maggots, Scoluding Brown Recluse Spider, Marbled Cellar Spider, Longbodied Cellar Spider, Shortbodied Cellar Spider, Burrowing Wolf Spider, Pirate Wolf Spider), Spider Mites, Springtails, Striped Grassworm, Jown Dog Tick and ticks that may transmit Lyme Disease), Two-lined Spittle Bugs, Webworms (includebworm, Grass Webworm, Tropical Sod Webworm), Weevils (including Hyperodes and Bluegrass) are table below, the Pre-Harvest Interval (PHI) refers to the minimum number of days that must pass be last application and the date of harvest. Reapply if insects return, but not more than once every seven or re-harvest interval.	Bronzed Cutworm, lafer, European Inbug, Ground Two-spotted irrobest, leafhoppers, cket, ay Beetle), on, Yellow Ground owbugs, Spiders Spider, Thinlegged Ticks (including ling Sod Ind Wireworms	awn ranules
ichokes et (Garden)	5 Days 1 Day	= .5
assica Vegetables: iccoli Raab (rapini), Chinese Cabbage (bok choy), Collards, Kale, Mizuna, Mustard Greens, Mustard Spinach, Rape (ad & Stem Brassica Vegetables: accoli, Chinese Broccoli (gai Ion, white flowering broccoli), Brussels Sprouts, Cauliflower, Cavalo Brocc		
hlrabi, Cabbage, Chinese Cabbage (napa) rn (sweet)	3 Days	
curbit Vegetables: ayote (fruit), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Gourd ilible (including hyotan, cucuzza, hechima, Chinese orkra), Mormordica spp. (includes balsam apple, bal ar, bitter melon, Chinese cucumber), Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenst elon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple m nta Claus melon, and snake melon), Pumpkin, Summer Squash (includes crookneck squash, scallop si aightneck squash, vegetable marrow, zucchini), Winter Squash (includes butternut squash, calabaza, bbard squash, acorn squash, and spaghetti squash), Watermelon (includes hybrids and varieties) ad Lettuce	lsam naw nelon, 3 Days	ects + ver
gume Vegetables - Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried Shelled Peas varf Pea, Edible-pod Pea, Snow Pea, Sugar Snap Pea, Pigeon Pea, English Pea, Garden Pea, Green Pea. gume Vegetables - Succulent Edible-Podded Beans, Succulent Shelled Beans, and Dried Shelled B nner Bean, Snap Bean, Wax Bean, Asparagus Bean, Chinese Longbean, Moth Bean, Yardlong Bean, skbean, Soybean (immature seed), Swordbean, Lima Bean, Broad Bean (Fava Bean), Blackeyed Pea, uthern Pea, Grain Lupin, Sweet Lupin, White Lupin, White Sweet Lupin, Field Bean, Kidney Bean, Navy Rean, Paray Rean, Adryki Rean, Cations Courses Crowder Pea Worth Rean, Kidney Bean, Ridey Bean, Navy Rean, Broad Rean, May Rean, Rides	s: (Succulent) , Lentil 21 Days (Dried) Beans: 1 Day (Succulent) Bean, 21 Days	
ito Bean, Tepary Bean, Adzuki Bean, Catjang, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Rice Bear an, Chickpea (Garbanzo Bean), Guar, Lablab Bean <mark>uiting Vegetables:</mark> pplant, Okra, Pepino (Melon pear), Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet		6 13499 01019 3
anuts ot and Tuber Vegetables (except Sugar Beets): acacha, Arrowroot, Garden Beet, Edible Burdock, Edible Canna, Carrot, Cassava (Bitter and Sweet), Celeriac elery Root), Chayote (Root), Turnip-Rooted Chervil, Chicory, Chufa, Dasheen (Taro), Ginger, Ginseng, Horse	radish, 21 Days	
ren, Turnip-Rooted Parsley, Parsnip, Oriental Radish (Daikón), Radish, Rutabaga, Śalsify (Oyster Plant), Blac sify, Spanish Salsify, Skirret, Sweet Potato, Tanier (Cocoyam), Turmeric, Turnip, Yam Bean, Yam (True) tato		

BERRIES, FRUITS, AND NUTS

Caneberries including: Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, Raspberries

Tree Nuts Group including:
Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia nut, Pistachio, Walnut (Black and English)



See Back Panel For Additional Precautionary Statements and Directions for Use.


Net Weight: 10 LB. (4.54 kg)

Active Ingredients:

Contains Peanut by-products

.....99.856%







GardenTech is a registered trademark of Gulfstream Home & Garden, Inc Sevin is a registered trademark of Tessenderlo Kerley, Inc. Scotts, Turf Builder and Edgeguard are registered trademarks of OMS Investments, Inc. Sta-Green is a registered trademark of LF, LLC.

QUICK FACTS

Protects edibles from insect damage Kills lawn-damaging insects that cause your lawn to thin and turn brown Can be used as an outdoor barrier treatment Do not allow people or pets to enter the treated area until all dusts have settled, watering-in is complete, an the treated area has dried

Fruit and Vegetable Gardens, Ornamental Plants and Shrubs, Flowers and Ground Covers, Around Structures, Lawns

Protects over 175 listed fruits and vegetables Tomatoes, Peppers, Zucchini, Sugar Snap Peas, Carrots, Blueberries, Grapes, Raspberries, Broccoli, Kale, Spinach, Cucumber, Pumpkin,

Kills Over 100+ Insects, as listed Ants, Aphids, Armyworms, Fleas, Ticks, Beetles, Cutworms, Japanese Beetles, Chinch Bugs, Caterpillars, and other listed lawn

I-4 lbs. per 1,000 square feet

• Call poison control center or doctor immediately for treatment advice wallowed
 Have person sip a glass of water if able to swallow.
 Do not induce vomiting unless told to do so by the poison control center or doctor. Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-420-9347 for Emergency Assistance. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatmet is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

For Information Regarding the Use of this Product Call 1-866-945-5033. PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals)

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

This pesticide is extremely toxic to fish, aquatic invertebrates, oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USEIt is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the following "Directions for Use" carefully and fully before applying this product. Apply this product directly to the lawn, home perimeter, or garden area. Water treated area as directed on this label. Do not water to the point of run-off.

Do not make applications during rain. Do apply when insect activity or damage is first observed in the lawn (lawn thins and turns brown or moths fly from the

grass when you mow)
Repeat application is necessary only if there are signs of renewed insect activity.
Do not allow people or pets to enter the treated area until all dusts have settled, watering-in is complete, and the treated area has dried. In New York State, this product may NOT be applied to any grass or turf area (lawns within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch))
In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application. Sevin® Insect Killer Lawn Granules is designed for use as a ready-to-use granular insecticide to control various adult and larvae above surfaces of lawns, and gardens. The active ingredients in this product have sufficient knockdown and residual activity to provide curative and residual control of surface feeding pests.

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347 If you have questions relating to product use or application, please call (866) 945-5033 If you have a product spill or transportation emergency call (800) 424-9300 For Consumer Refund: TechPac, LLC PO Box 95437 Palatine, IL 60095-0437

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry area. Always store pesticides in the original container. Store away from food and pet food.

Pesticide Disposal and Container Handling

Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent permitted by applicable law, User agrees to hold Seller harmless for any claims

LAWN, PERIMETER, and GARDEN APPLICATIONS Broadcast **Sevin® Insect Killer Lawn Granules** to provide uniform coverage over the treatment area. Locate the listed pest to be controlled in the following table and follow the specified application rate to provide excellent control under typical conditions. When treating for multiple pests, apply the higher of the specified rates. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity. Repeat application no more than once per 7 days. Do not exceed 4 lb per 1,000 ft² per application.

Best results are obtained if the treated area is thoroughly watered immediately after application. Residential Lawn Application: Broadcast this product with spreader equipment to ensure uniform coverage over the treatment area. Perimeter Application: Apply to ground area 5 to 10 feet wide around and adjacent to the structure. Ornamental & Flower Garden Application: Use this product as a spot treatment or area treatment for pest infestations on or around ornamental trees, shrubs, flowers and ground covers. Certain plants such as annuals may be sensitive to the watered-in

granules. A limited area should be treated and observed for one week prior to application to the entire planting. Fruit & Vegetable Garden Application: Apply this product to control listed pests in home fruit and vegetable gardens either as a preventative measure or to control Preventative Applications:

This product may be applied just prior to planting or after plants emerge and worked into the top 4 to 6 inches of soil. Use 1-4 lb of product for a 1,000 sq ft area and immediately water the area thoroughly after application for best results. Applications to Control Existing Infestations: Apply this product to the garden area using 1 to 4 lb of product for a 1,000 sq ft area. Best results are obtained if the treated area is thoroughly watered immediately after application. Treat when insects first appear and retreat as necessary.

SPREADER EQUIPMENT SETTINGS
Broadcast application should be performed with a calibrated broadcast spreader to evenly distribute the granules.

Approximate settings for several models are provided in the table below. Mow lawn first for best results.

SPREADER SETTINGS

			NOTE: Spreader settings are approximate. It is		
Broadcast Spreaders	Rate (lbs/1,000 ft ²)	Setting	recommended that you calibrate your spreader before		
Scotts® Turf Builder® Edgeguard® DLX	1 2 4	4 1/2 7 12	application since manufacturer's settings can vary among the same model of spreader. Broadcast Spreader : If your lawn is rectangular, apply two		
Scotts® Turf Builder® Edgeguard® Mini	1 2 4	3 3/4 5 1/2 12	border strips across each end for a turning area. If your lawn is an irregular shape, apply a border strip completely around it. Then apply product back and forth in the longesi		
Sta-Green® Rotary	1 2 4	3 4 3/4 9	direction. Turn the spreader off and on at the edge of each strip to prevent uneven application. Overlap the throw pattern by about a foot to ensure complete coverage.		
Drop Spreaders	Rate (lbs/1,000 ft²)	Setting	Drop Spreader: Apply granules with spreader going in the same direction as the longest part of the lawn. Apply two (2)		
Scotts® Turf Builder® Edgeguard® Classic	1 2 4	5 1/4 8 10	header strips first. To ensure thorough coverage, make sure to overlap wheel marks. Shutoff the spreader when finished or when the header strips have been reached.		



TechPac, L.L.C. 1000 Parkwood Circle, Suite 700 Atlanta, GA, 30339 EPA Reg. No. 279-3346-71004 13499 01019 3 EPA Est. 67572-GA-1_A, 67572-GA-2_B

Subscript used is first letter of run code on container.

www.gardentech.com